

CERTAIN SUBCLASSES OF CONVEX FUNCTIONS WITH POSITIVE AND MISSING COEFFICIENTS BY USING A FIXED POINT

SH. NAJAFZADEH, M. ESHAGHI GORDJI AND A. EBADIAN

ABSTRACT. By considering a fixed point in unit disk Δ , a new class of univalent convex functions is defined. Coefficient inequalities, integral operator and extreme points of this class are obtained.

1. INTRODUCTION

Let w be fixed point in $\Delta = \{z : |z| < 1\}$ and

$$A(w) = \{f \in H(\Delta) : f(w) = f'(w) - 1 = 0\}.$$

Let $M_w = \{f \in A(w) : f \text{ is Univalent in } \Delta\}$, and M_w denoted the subclass of $A(w)$ consist of the functions of the form

$$(1.1) \quad f(z) = \frac{1}{z-w} + \sum_{n=k}^{+\infty} a_n(z-w)^n, a_n \geq 0, z \neq w$$

The function $f(z)$ in M_w is said to be Convex of order η ($0 < \eta < 1$) if and only if

$$Re\left\{1 + \frac{(z-w)f''(z)}{f'(z)}\right\} > \eta, \quad (z \in \Delta).$$

The similar definitions for uniformly convex and starlike functions are introduced by Goodman in [4]. For the function $f(z)$ in M_w , Ghanim and Darus have defined an operator as follows:

$$\begin{aligned} I^0 f(z) &= f(z) \\ I^1 f(z) &= (z-w)f'(z) + \frac{2}{z-w} \\ I^2 f(z) &= (z-w)(I^1 f(z))' + \frac{2}{z-w} \end{aligned}$$

and for $m = 1, 2, \dots$

$$\begin{aligned} (1.2) \quad I^m f(z) &= (z-w)(I^{m-1} f(z))' + \frac{2}{z-w} \\ &= \frac{1}{z-w} + \sum_{n=k}^{+\infty} n^m a_n (z-w)^n. \end{aligned}$$

We note that various other subclasses have been studied rather extensively by many authors. (see also [3] and [6])

2000 *Mathematics Subject Classification.* 30C45, 30C50.

Key words and phrases. Subordination; P-valent function; Coefficient estimate; Fixed point; Distortion bound and Radii of starlikeness and Convexity.

Definition 1.1. A function $f(z)$ belonging to the class M_w is in the class $M_w(A, B, m)$ if it satisfies the condition

$$(1.3) \quad \left| \frac{\frac{1}{2}[(z-w)\frac{(I^m f(z))''}{(I^m f(z))'} + 1] + \frac{1}{2}}{\frac{B}{2}[(z-w)\frac{(I^m f(z))''}{(I^m f(z))'} + 1] + \frac{A+1}{2}} \right| \leq 1$$

for some $-1 \leq B < A < 1$, $0 \leq A \leq 1$.

2. MAIN RESULTS

First we obtain coefficient inequalities for functions in $M_w(A, B, m)$. Then we prove the linear combination property.

Theorem 2.1. *The function $f(z)$ of the form (1.1) belongs to $M_w(A, B, m)$ if and only if*

$$(2.1) \quad \sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] a_n \leq 1 + A - B.$$

Proof. Suppose (2.1) holds and

$$(2.2) \quad H = \left| \frac{1}{2}(z-w)(I^m f(z))'' + (I^m f(z))' \right| - \left| \frac{B}{2}(z-w)(I^m f(z))'' + \frac{B+A}{2} \right|$$

Replacing $(I^m f(z))'$ and $(I^m f(z))''$ by their series expansion for $0 < |z-w| = r < 1$, we have

$$\begin{aligned} H &= \left| \sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (n+1) a_n (z-w)^{n-1} \right| - \left| \frac{1}{2} (B-A-1) \frac{1}{(z-w)^2} + \sum_{n=k}^{+\infty} n^{m+1} \left[\frac{1}{2} (Bn+A+1) \right] a_n (z-w)^{n-1} \right| \\ &\leq \left| \sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (n+1) a_n r^{n-1} - \frac{1}{2} (1+A-B) \frac{1}{r^2} + \sum_{n=k}^{+\infty} n^{m+1} \left[\frac{1}{2} (Bn+A+1) \right] a_n r^{n-1} \right| \end{aligned}$$

Since this inequality holds for all r ($0 < r < 1$), making $r \rightarrow 1$, we have

$$H \leq \sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} [n(B+1) + A + 2] a_n - \frac{1}{2} (1+A-B)$$

by (2.1)

$$H \leq 0.$$

So we have the required result.

Conversely, let

$f(z) = \frac{1}{z-w} + \sum_{n=k}^{+\infty} a_n (z-w)^n$ and (1.3) holds, so we have

$$\left| \frac{\sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (n+1) a_n (z-w)^{n-1}}{\frac{1}{2} (B-A-1) \frac{1}{(z-w)^2} + \sum_{n=k}^{+\infty} n^{m+1} \left[\frac{1}{2} (Bn+A+1) \right] a_n (z-w)^{n-1}} \right| \leq 1,$$

or

$$\left| \frac{\sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (n+1) a_n (z-w)^{n+1}}{\frac{1}{2} (1+A+B) - \sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (Bn+A+1) a_n (z-w)^{n+1}} \right| \leq 1$$

Since for all $z, \operatorname{Re}(z) \leq |z|$, so

$$\operatorname{Re} \left\{ \frac{\sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (n+1) a_n (z-w)^{n+1}}{\frac{1}{2} (1+A+B) - \sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (Bn+A+1) a_n (z-w)^{n+1}} \right\} \leq 1.$$

Choosing $z - w = r$ with $0 < r < 1$, we get

$$(2.3) \quad \frac{\sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (n+1) a_n r^{n+1}}{\frac{1}{2} (1+A-B) - \sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (Bn+A+1) a_n r^{n+1}} \leq 1.$$

Upon clearing the denominator in (2.3) and letting $r \rightarrow 1$, we get

$$\sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (n+1) a_n \leq \frac{1}{2} (1+A-B) - \sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} (Bn+A+1) a_n,$$

or

$$\sum_{n=k}^{+\infty} \frac{1}{2} n^{m+1} [n(B+1) + A + 2] a_n \leq \frac{1}{2} (1+A-B).$$

Hence the proof is complete. $\square$

Theorem 2.2. Let $f_j(z)$ defined by

$$(2.4) \quad f_j(z) = \frac{1}{z-w} + \sum_{n=k}^{+\infty} a_{n,j} (z-w)^n, j = 1, 2, \dots$$

be in the class $M_w(A, B, m)$, then the function

$$F(z) = \sum_{j=1}^t d_j f_j(z) \quad , \quad d_j \geq 0$$

is also in $M_w(A, B, m)$, where $\sum_{j=1}^t d_j = 1$.

Proof. Since $f_j(z) \in M_w(A, B, m)$, by (2.1) we have,

$$(2.5) \quad \sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] a_{n,j} \leq 1 + A - B \quad , j = 1, 2, \dots$$

also

$$\begin{aligned} F(z) &= \sum_{j=1}^t d_j \left(\frac{1}{z-w} + \sum_{n=k}^{+\infty} a_{n,j} (z-w)^n \right) \\ &= \frac{1}{z-w} \sum_{j=1}^t d_j + \sum_{n=k}^{+\infty} \left(\sum_{j=1}^t d_j a_{n,j} \right) (z-w)^n \\ &= \frac{1}{z-w} + \sum_{n=k}^{+\infty} s_n (z-w)^n \quad \text{where} \quad s_n = \sum_{j=1}^t d_j a_{n,j} \end{aligned}$$

But

$$\begin{aligned} \sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] s_n &= \sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] \left[\sum_{j=1}^t d_j a_{n,j} \right] \\ &= \sum_{j=1}^t d_j \left\{ \sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] a_{n,j} \right\} \end{aligned}$$

by (2.5)

$$\leq \sum_{j=1}^t d_j (1 + A - B) = 1 + A - B.$$

Now the proof is complete. $\square$

3. EXTREME POINTS AND INTEGRAL OPERATORS

In the last section we investigate about extreme points of $M_w(A, B, m)$ and verify the effect of two operators on functions in the class $M_w(A, B, m)$.

Theorem 3.1. *Let*

$$f_0(z) = \frac{1}{z-w}$$

and

$$(3.1) \quad f_n(z) = \frac{1}{z-w} + \frac{1+A-B}{n^{m+1}[n(B+1)+A+2]}(z-w)^n, \quad n \geq k$$

then $f(z) \in M_w(A, B, m)$ if and only if it can be expressed in the form $f(z) = \sum_{n=1}^{\infty} C_n f_n(z)$ where $C_n \geq 0$, $C_i = 0$ ($i = 1, 2, \dots, k-1$), $\sum_0^{+\infty} C_n = 1$

Proof. Let $f(z) = \sum_0^{+\infty} C_n f_n(z)$, so,

$$\begin{aligned} f(z) &= \frac{C_0}{z-w} + \sum_{n=1}^{k-1} C_n \left[\frac{1}{z-w} + \frac{1+A-B}{n^{m+1}[n(B+1)+A+2]}(z-w)^n \right] + \\ &\quad \sum_{n=k}^{+\infty} C_n \left[\frac{1}{z-w} + \frac{1+A-B}{n^{m+1}[n(B+1)+A+2]}(z-w)^n \right] \\ &= \frac{1}{z-w} + \sum_{n=k}^{+\infty} \frac{1+A-B}{n^{m+1}[n(B+1)+A+2]} C_n (z-w)^n. \end{aligned}$$

Since

$$\begin{aligned} &\sum_{n=k}^{+\infty} \frac{1+A-B}{n^{m+1}[n(B+1)+A+2]} \frac{n^{m+1}[n(B+1)+A+2]}{1+A-B} C_n \\ &= \sum_{n=k}^{+\infty} C_n = \sum_{n=1}^{+\infty} C_n = 1 - C_0 \leq 1, \end{aligned}$$

so $f(z) \in M_w(A, B, m)$.

Conversely, suppose that $f(z) \in M_w(A, B, m)$. Then by (2.1) we have

$$0 \leq a_n \leq \frac{1+A-B}{n^{m+1}[n(B+1)+A+2]}.$$

By setting

$$C_n = \frac{n^{m+1}[n(B+1)+A+2]}{1+A-B} a_n, \quad n \geq 1$$

$C_i = 0$ ($i = 1, 2, \dots, k-1$), $C_0 = 1 - \sum_{n=1}^{+\infty} C_n$, we obtain the required result. $\square$

Theorem 3.2. *Let γ be a real number such that $\gamma > 1$. If $f(z) \in M_w(A, B, m)$, then the functions*

$$H_1(z) = \frac{\gamma-1}{(z-w)^\gamma} \int_w^z (t-w)^{\gamma-1} f(t) dt \quad \text{and}$$

$$H_2(z) = C \int_0^1 \nu^C f(\nu(z-w) + w) d\nu, \quad C \geq 1$$

are also in the same class.

Proof. Let $f(z) \in M_w(A, B, m)$, then a simple calculation shows that,

$$H_1(z) = \frac{1}{z-w} + \sum_{n=k}^{+\infty} \frac{1}{\gamma+n} a_n (z-w)^n$$

$$H_2(z) = \frac{1}{z-w} + \sum_{n=k}^{+\infty} \frac{C}{C+n+1} a_n (z-w)^n$$

Since $\frac{C}{C+n+1} < 1$ and $\frac{1}{\gamma+n} < 1$, by using Theorem 2.1, we get,

$$\sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] a_n \frac{1}{\gamma+n} < \sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] a_n < 1 + A - B$$

and

$$\sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] a_n \frac{C}{C+n+1} < \sum_{n=k}^{+\infty} n^{m+1} [n(B+1) + A + 2] a_n < 1 + A - B$$

Hence by Theorem (2.1) we conclude that $H_1(z)$ and $H_2(z)$ are in the class $M_w(A, B, m)$. So the proof is complete. $\square$

REFERENCES

- [1] F. Ghanim and M. Darus, On new subclass of analytic univalent function with negative coefficient I, Int. J. Contemp. Math. Sciences, vol. 3. No27, (2008), 1317 – 1329.
- [2] ———, Some subordination results associated with certain subclass of analytic meromorphic function, J. Math. and statistics, 4(2), (2008), 112 – 116.
- [3] F. Ghanim, M. Darus and S. Sivasubramanian, On subclass of analytic univalent function, Int. J. pure Appl., 40, (2007), 307 – 319.
- [4] A .W. Goodman, On uniformly starlike functions, J. Math. Anal. Appl, 155, (1991), 364–370.
- [5] ———, On uniformaly convex functions, Ann. polon. Math, 56(1), (1991), 87–92.
- [6] Sh. Najafzadeh, A. Tehranchi and S. R. Kulkarni, Application of differential operator on p-valent meromorphic functions, Anal. Univ. oradea, Fasc. Math. Tom XII, (2005), 75 – 90.

Sh. Najafzadeh

Department of Mathematics, Faculty of Science,
University of Maragheh, Maragheh, Iran
E.mail: najafzadeh1234@yahoo.ie

M. Eshaghi Gordji Department of Mathematics, Semnan University,
P. O. Box 35195-363, Semnan, Iran
E.Mail: madjid.eshaghi@gmail.com

A. Ebadian

Department of Mathematics, Faculty of Science,
Urmia University , Urmia, Iran
E.mail: aebadian@yahoo.com